



Enzyme Research Laboratories

1801 Commerce Drive
South Bend, IN U.S.A. 46628
(574) 288-2268 * Fax (574) 288-2272
Website: www.enzymeresearch.com
Email: info@enzymeresearch.com

Murine Anti-Factor X

Clone 509

Factor X (Mr 59,000) is a vitamin K-dependent plasma protein zymogen that plays a central role as the substrate for both the intrinsic (factor VIIa, tissue factor) and extrinsic (factor IXa, factor VIIIa) pathways. In the presence of cofactor factor Va, phospholipid, and Ca^{2+} , activated factor X cleaves two peptide bonds in prothrombin to form thrombin. Antibody binds human factor X in solid-phase ELISA and Western blot.

Description

Antibody Source: mouse monoclonal, IgG_{2a}

Antigen Species Bound: human

Specificity: factor X light chain

Immunogen: human factor X

Formulation and Storage

Purity: Purified by protein G affinity chromatography from serum-free cell culture supernatant.

Product Formulation: Lyophilized from a ≥ 1 mg/ml solution in 20 mM NaH_2PO_4 0.15 M NaCl, 1.0% (w/v) mannitol, pH 7.4. Concentration determined by absorbance measurement at 280 nm and using an extinction coefficient of 1.4 ($\epsilon_{0.1\%}$).

Reconstitution: Reconstitute with deionized water.

Storage: Store lyophilized or reconstituted and aliquoted material at $-20^\circ C$ for prolonged periods. Avoid freeze-thaw cycles. Alternatively, add 0.02% (w/v) sodium azide to reconstituted solution and store at $4^\circ C$.

Country of Origin: USA

Size Options: 0.1 mg or 0.5 mg

Applications

Working Concentration: Approximately 1-5 $\mu g/ml$. Researcher should titer antibody in specific assay.

ELISA: Binds human factor X and Xa, and can be used in sandwich ELISA with Mab HFX.

Immunoblotting: Binds human factor X light chain, under reduced conditions and non-reduced conditions.

