



Enzyme Research Laboratories

1801 Commerce Drive
South Bend, IN U.S.A. 46628
(574) 288-2268 * Fax (574) 288-2272
Website: www.enzymeresearch.com
Email: info@enzymeresearch.com

Murine Anti-Plasminogen

Clone 030

Plasminogen, precursor of the active protease plasmin, is a single chain glycoprotein of 92 kDa. Found in plasma at a concentration of 200 ug/ml, it contains 5 disulfide-bonded structures termed "kringles" and a serine protease domain at the carboxy-terminus. Plasmin is primarily responsible for digesting fibrin clots. Mab HPG-030 binds human plasminogen kringle 5 B-chain in solid-phase ELISA. Antibody inhibits plasminogen activation by streptokinase, tPA and uPA.

Description

Antibody Source: mouse monoclonal, IgG₁

Antigen Species Bound: human

Specificity: Kringle 5 B-chain

Immunogen: human plasminogen

Formulation and Storage

Purity: Purified by protein G affinity chromatography from serum-free cell culture supernatant.

Product Formulation: Lyophilized from a ≥ 1 mg/ml solution in 20 mM NaH₂PO₄ 0.15 M NaCl, 1.0% (w/v) mannitol, pH 7.4. Concentration determined by absorbance measurement at 280 nm and using an extinction coefficient of 1.4 ($\epsilon_{0.1\%}$).

Reconstitution: Reconstitute with deionized water.

Storage: Store lyophilized or reconstituted and aliquoted material at -20°C for prolonged periods. Avoid freeze-thaw cycles. Alternatively, add 0.02% (w/v) sodium azide to reconstituted solution and store at 4°C.

Country of Origin: USA

Size Options: 0.1 mg or 0.5 mg

Applications

Working Concentration: Approximately 1-5 μ g/ml. Researcher should titer antibody in specific assay.

ELISA: Binds plasminogen kringle 5 B-chain.

Inhibition: Inhibits plasminogen activation by streptokinase, tPA and uPA.