

Website: www.enzymeresearch.com Email: info@enzymeresearch.com

Murine Anti-Plasminogen

Clone 039


Size Options:

Global Coag

Plasminogen, precursor of the active protease plasmin, is a single chain glycoprotein of 92 kDa. Found in plasma at a concentration of 200 ug/ml, it contains 5 disulfide-bonded structures termed "kringles" and a serine protease domain at the carboxy-terminus. Plasmin is primarily responsible for digesting fibrin clots. Mab HPG binds human plasminogen and kringle 5 by ELISA and western blots.

Description			
Antibody Source:		mouse monoclonal, IgG ₁	
Antigen Species Bound:		human	
Specificity:		kringle 5-B chain	
Immunogen:		human plasminogen	
Formulation and Storage			
Purity:	Purified by protein G affinity chromatography from serum-free cell culture supernatant.		
Product Formulation:	Lyophilized from a ≥ 1 mg/ml solution in 20 mM NaH ₂ PO ₄ 0.15 M NaCl, 1.0% (w/v) mannitol, pH 7.4. Concentration determined by absorbance measurement at 280 nm and using an extinction coefficient of 1.4 ($\epsilon_{0.1\%}$).		
Reconstitution:	Reconstitute with deionized water.		
Storage:	Store lyophilized or reconstituted and aliquoted material at -20°C for prolonged periods. Avoid freeze-thaw cycles. Alternatively, add 0.02% (w/v) sodium azide to reconstituted solution and store at 4°C.		
Country of Origin:	USA		

Applications		
Working Concentration:	Approximately 1-5 µg/ml. Researcher should titer antibody in specific assay.	
ELISA:	Binds plasminogen and kringle 5.	
Immunoblotting:	Binds plasminogen, at an approximate molecular weight of 88 kD, under non-reduced conditions.	



References

[1] D.J Davidson, C. Haskell, S. Majest, A. Kherzai, D.A. Egan, K.A. Walter, A. Schneider, E. Gubbins, L. Solomon, Z. Chen, R. Lesniewski, J. Henkin. Kringle 5 of Human Plasminogen Induces Apoptosis of Endothelial and Tumor Cells through Surface-Expressed Gluclose-Regulated Protein 78. (2005). Cancer Res. 65: 4663-4666.

0.1 mg or 0.5 mg